Kansas Wind & Prairie Untilled Areas - the Tallgrass Prairie Ecosystem Jewell Republic Washington Marshall Nemaha Brown Doniphan Atchison Jacksor Cloud Clay Mitchell Jefferson -eavenworth Ottawa Lincoln Dickinson Douglas Saline Osage Ellsworth Franklin Mario McPherson Rice Harvey Reno Allen Sedgwick Neosho Kingman Labette Montgomery 25 50 Miles Map compilation by the Kansas Geological Survey (KGS), University of Kansas, December, 2003. 1:1,500,000 Data sources: Base map (county boundaries) - TIGER 2000, US Census Bureau Untilled areas, The Nature Conservancy Untilled fragments, The Nature Conservancy Legend Counties Untilled areas "Untilled areas were identified via an interpretation of Landsat Thematic Mapper (TM) satellite imagery. These areas were classified as one of two types: 1) areas with largely intact natural or semi-natural vegetation, and 2) fragmented areas with a concentration of natural community remnants." (The Nature Untilled fragments Citation: The Nature Conservancy, 2003, Relatively Intact or "Untilled" Natural Community Landscapes (e.g. prairies, savannas) in the Osage Plains/Flint Hills Ecoregion, The Nature Conservancy. Conservancy, 2003)